

South West HP Operations 3rd Floor, 2 Rivergate Temple Quay, Bristol, BS1 6EH T +44 (0)300 303 8162

Follaton House, Plymouth Road, Totnes, Devon, TQ9 5NE

www.gov.uk/ukhsa

6/5/22

Dear Parent/Guardian,

I am writing to let you know that a number of children and staff at Brentry Primary School have had diarrhoea and/or vomiting over the last few days.

I have been advised by the South West Health Protection Team (UKHSA) that this is generally a self-limiting illness probably caused by a virus, but that it is very infectious. The most usual symptom is vomiting. Some people may have diarrhoea and/or abdominal pain. Symptoms rarely last for more than 24 – 48 hours. The incubation period (time it takes for the illness to develop) is between 12 and 48 hours.

If your child is affected, please keep them off school. They should not return until 48 hours after the diarrhoea and vomiting have stopped.

The virus is easily spread from person to person. Good hygiene by everyone in the family reduces the risk. This means washing hands with soap and warm water after going to the toilet, and before preparing or eating food as a minimum.

It is important to clean up carefully when someone has been sick as vomit is very infectious. It is important to clear up spills of vomit or faeces immediately, by thorough washing of the contaminated environment with detergent and hot water. For hard surfaces, (floor, work tops etc. an additional disinfection with a dilute solution of Milton or household bleach (according to manufacturer's instructions) will reduce the contamination. DO NOT MIX these substances with soap and water. Be sure that each family member uses separate towels and flannels which are changed and washed frequently.

Staying off school until fully recovered for 48 hours and hand washing are the most important elements of reducing the risk of infection.

We are ensuring the school is thoroughly cleaned to reduce any further risk.

If you need any further advice you can phone NHS 111. For more information on symptoms and management of viral gastro-enteritis please see https://www.nhs.uk/conditions/diarrhoea-and-vomiting/

Yours faithfully

Mr Clarke